


LAC OPERON

**BY:NEHA MEGHANI
DEPARTMENT OFBIOCHEMISTRY**



Operons

- An operon is a group of genes that are **transcribed at the same time**.
- They usually control an important biochemical process.
- They are **only found in prokaryotes**.



[Jacob, Monod & Lwoff](#)

The *lac* Operon

- The lac operon consists of **three genes** each involved in processing the sugar lactose
- One of them is the gene for the enzyme **β -galactosidase**
- This enzyme hydrolyses lactose into glucose and galactose.

Adapting to the environment


- *E. coli* can use either glucose, which is a monosaccharide, or lactose, which is a disaccharide
- However, lactose needs to be hydrolysed (digested) first
- So the bacterium prefers to use glucose when it can.

Four situations are possible

1. When glucose is **present** and lactose is **absent** the E. coli does **not** produce β -galactosidase.
2. When glucose is **present** and lactose is **present** the E. coli does **not** produce β -galactosidase.
3. When glucose is **absent** and lactose is **absent** the E. coli does **not** produce β -galactosidase.
4. When glucose is **absent** and lactose is **present** the E. coli **does** produce β -galactosidase.


1. When lactose is absent

- A repressor protein is continuously synthesised. It sits on a sequence of DNA just in front of the *lac* operon, the **Operator site**
- The **repressor protein** blocks the **Promoter site** where the RNA polymerase settles before it starts transcribing



2. When lactose is present


- A small amount of a sugar allolactose is formed within the bacterial cell. This fits onto the repressor protein at another active site (**allosteric site**)
- This causes the repressor protein to change its shape (a **conformational change**). It can no longer sit on the operator site. RNA polymerase can now reach its promoter site



3. When both glucose and lactose are present


- This explains how the *lac* operon is transcribed only when lactose is present
- BUT..... this does **not** explain why the operon is not transcribed when **both** glucose and lactose are present.

- When glucose and lactose are present RNA polymerase can sit on the promoter site but it is unstable and it keeps falling off



4. When glucose is absent and lactose is present

- Another protein is needed, an **activator protein**. This stabilises RNA polymerase.
- The activator protein only works when glucose is absent
- In this way *E. coli* only makes enzymes to metabolise other sugars in the absence of glucose.



Summary

Carbohydrates	Activator protein	Repressor protein	RNA polymerase	<i>lac</i> Operon
+ GLUCOSE + LACTOSE	Not bound to DNA	Lifted off operator site	Keeps falling off promoter site	No transcription
+ GLUCOSE - LACTOSE	Not bound to DNA	Bound to operator site	Blocked by the repressor	No transcription
- GLUCOSE - LACTOSE	Bound to DNA	Bound to operator site	Blocked by the repressor	No transcription
- GLUCOSE + LACTOSE	Bound to DNA	Lifted off operator site	Sits on the promoter site	Transcription

Thank You

